

Knowledge of and attitudes toward epilepsy surgery among neurologists in Saudi Arabia

Bandar Aljafen, MD, Majed Alomar, MBBS, Nawaf Abohamra, MBBS, Mohammed Alanazy, MD, Fawaz Al-hussain, MD, Ziad Alhumayyd, MD, Yousef Mohammad, MD, Taim Muayqil, MD.

ABSTRACT

الأهداف: دراسة مستويات المعرفة والمواقف تجاه جراحة الصرع بين أطباء الأعصاب في المملكة العربية السعودية وتقييم العوامل التي تؤثر على معرفة الأطباء ومواقفهم

المنهجية: دراسة مقطعية رصدية كمية في مدينة جامعة الملك سعود الطبية، الرياض. تم جمع البيانات باستخدام استبيان حديث ومتطور ذاتي الإدارة عبر الإنترنت. تضمن الاستبيان 3 أقسام: المعلومات الديموغرافية، والمعرفة، والمواقف التي أرسلت بعد ذلك إلى أطباء الأعصاب في المملكة العربية السعودية من ديسمبر 2016م إلى مارس 2017م.

النتائج: توافق 106 من أطباء الأعصاب مع معايير الاشتغال لدينا. ثمانون في المائة من المشاركين لديهم مركز واحد على الأقل من مراكز جراحة الصرع في مدينتهم، وأشار 78% إلى أن لديهم القدرة على الوصول إلى المراكز ذات الخبرة والموارد الكافية لتمكين الاختيار المناسب للمرضى المرشحين لجراحة الصرع. كان 57.5% فقط من أطباء الأعصاب لديهم مستوى كاف من المعرفة بشأن جراحة الصرع وقام أطباء الأعصاب ذوي المستوى الأعلى من المعرفة بإحالة المزيد من المرضى إلى وحدات مراقبة الصرع وناقشوا جراحة الصرع في كثير من الأحيان مع مرضاهم. بشكل عام، كان لدى أكثر من نصف أطباء الأعصاب (52.8%) موقف إيجابي تجاه جراحة الصرع وكان هناك ارتباط إيجابي كبير بين درجات المعرفة والمواقف ($p < 0.001$).

الخلاصة: يبدو أن أطباء الأعصاب في المملكة العربية السعودية لديهم معرفة متوسطة ومواقف إيجابية تجاه جراحة الصرع. أثر على معرفتهم عوامل كثيرة منها مكان آخر شهادة في علم الأعصاب، نوع المستشفى الذي ينتمي له الممارس وسهولة الوصول إلى مراكز الصرع ذات الخبرة والموارد وارتبطت المعرفة الكافية بشكل إيجابي مع المواقف

Objectives: To investigate the levels of knowledge and attitudes toward epilepsy surgery among neurologists in Saudi Arabia and evaluate the factors that affect the physicians' knowledge and attitudes.

Methods: A quantitative observational cross-sectional study conducted at King Saud University Medical

City, Riyadh. The data were collected using a newly developed, self-administered online questionnaire. The questionnaire contained 3 sections: demographic information, knowledge, and attitudes which then sent to neurologist in Saudi Arabia from December 2016 to March 2017.

Results: A total of 106 neurologists met our inclusion criteria. Eighty percent of the participants had at least one epilepsy center in their city, and 78% indicated that they had access to adequate expertise and resources to enable the appropriate selection of epilepsy surgical candidates. Only 57.5% of the neurologists had a sufficient level of knowledge regarding epilepsy surgery. Neurologists with higher level of knowledge referred more patients to EMU and discussed epilepsy surgery more often with their patients. Overall, more than half of the neurologists (52.8%) had a positive attitude toward epilepsy surgery. There was a significantly positive correlation between the scores of knowledge and attitude ($p < 0.001$).

Conclusion: Neurologists in Saudi Arabia appear to have moderate knowledge of and positive attitudes toward epilepsy surgery. The place of the last neurology certificate, type of practicing hospital, and access to expertise and resources, affected their knowledge. Adequate knowledge was positively correlated with attitude.

*Neurosciences 2020; Vol. 25 (1): 43-49
doi: 10.17712/nsj.2020.1.20190051*

From the Division of Neurology (Aljafen, Alanazy, Al-hussain, Alhumayyd, Mohammad, Muayqil), Department of Internal Medicine, King Saud University Medical City and College of Medicine, King Saud University, and from National Neurosciences Institute (Alomar, Abohamra), King Fahad Medical City, Riyadh, Kingdom of Saudi Arabia.

Received 19th April 2019. Accepted 26th September 2019.

*Address correspondence and reprint request to: Dr. Bandar N. Aljafen, College of Medicine, King Saud University, Riyadh, Kingdom of Saudi Arabia. E-mail: bandaraljafen@gmail.com
ORCID ID: <https://orcid.org/0000-0001-8845-4020>*

Epilepsy is one of the most common neurological disorders in Saudi Arabia, with an estimated prevalence of 6.54 per 1000 individuals.¹ It is estimated that 20-40% of newly diagnosed patients are expected to fail seizure control with medical treatment.² Refractory epilepsy is commonly defined as the failure to achieve freedom from seizures despite two or more antiepileptic drugs (AEDs).³ Patients with drug-resistant epilepsy (DRE) have a lower quality of life and face more social difficulties than patients with controlled epilepsy.⁴⁻⁶ More than half of the economic burden of epilepsy is accounted for by patients with refractory epilepsy.⁷ Epilepsy surgery is the best evidence-based treatment option for these patients with drug resistant focal epilepsy.^{8,9} The current guidelines for patient referral to epilepsy surgery were published in 2003 by the American Academy of Neurology.⁹ The benefits of epilepsy surgery may not only reduce the frequency of seizures but might also lower mortality and improve the quality of life.¹⁰ Despite its proven efficacy and favorable outcome, the average referral delay of an epilepsy surgery candidate is more than 20 years.¹¹ This delay is predominately attributed to non-adherence to referral guidelines, which results from a lack of awareness and familiarity with them.¹² Another factor that contributes to the delay is patients' misbeliefs regarding the surgery's risks and benefits, which are highly influenced by physicians' lack of knowledge.¹³ Epilepsy surgery was introduced to Saudi Arabia in 1998.¹⁴ Several epilepsy surgery centers have subsequently been established across the country. A survey in 2013 showed that 56% of health-care professionals in Saudi Arabia were not aware that surgery was a treatment option for patients with epilepsy.¹⁵ Our aims in this study were to investigate the levels of knowledge and attitudes toward epilepsy surgery among neurologists in Saudi Arabia and evaluate the factors that affect the physicians' knowledge and attitudes.

Methods. A quantitative observational cross-sectional study was conducted at King Saud University Medical City (KSUMC), Riyadh, Kingdom of Saudi Arabia. Both adult and pediatric neurologists in Saudi Arabia were included in the study population. Institutional review board approval was granted prior

to data collection. The questionnaire was then sent to adult and pediatric neurologist in Saudi Arabia from December 2016 to March 2017.

Questionnaire. An anonymous, self-administered online questionnaire was completed using the Google Documents platform. The questionnaire was developed by 3 neurologists (BJ, TM, and MHA) based on other studies that have measured knowledge, attitudes and perceptions towards epilepsy surgery.¹⁶⁻¹⁹ The questionnaire contained 3 sections: demographic information, knowledge, and attitudes. The demographic section included age, gender, training site (residency or fellowship training program), subspecialty, experience, and other questions regarding access and referral habits for epilepsy surgery and epilepsy monitoring unit (EMU). The knowledge section included 11 items that evaluated different aspects of epilepsy surgery knowledge. Each item had one correct answer, one or more incorrect answers, and an "I don't know" option. A correct response was awarded one point, while any other response was awarded none. Individuals who scored 7 or more were considered to have sufficient knowledge. The attitude section contained 8 items, including both positive and negative attitude statements. These items used a 5-point Likert

Table 1 - Responses to the demographics section.

| Variables | n (%) |
|---------------------------------------------------------|-----------|
| <i>Gender</i> | |
| Male | 75 (70.8) |
| Female | 31 (29.2) |
| <i>Age (Years)</i> | |
| <40 | 52 (49.1) |
| 40-50 | 41 (38.7) |
| >50 | 13 (12.2) |
| <i>Place of the most recent neurology certification</i> | |
| North America | 52 (49.1) |
| Saudi Arabia | 44 (41.5) |
| Other | 10 (9.4) |
| <i>Region of practice</i> | |
| Central | 59 (55.7) |
| Western | 31 (29.2) |
| Eastern | 12 (11.3) |
| Southern | 2 (1.9) |
| Northern | 2 (1.9) |
| <i>Type of practicing institute</i> | |
| Ministry of health hospitals | 38 (35.8) |
| University hospitals | 31 (29.2) |
| Military hospitals | 24 (22.6) |
| Other | 13 (12.3) |

Disclosure. Authors have no conflict of interests, and the work was not supported or funded by any drug company.

Table 2 - Responses to the knowledge section.

| Statements | n (%) |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|
| <i>What is the approximate percentage of patients who experience clinically significant (i.e., disabling) and permanent adverse effects after anterior temporal lobectomy?</i> | |
| <5% (correct answer) | 38 (35.8) |
| 5-10% | 33 (31.1) |
| 11-50% | 3 (2.8) |
| >50% | 1 (0.9) |
| I don't know | 31 (29.2) |
| <i>Is there a general agreement on the definition of drug-resistant epilepsy?</i> | |
| Yes (correct answer) | 79 (74.5) |
| No / I don't know | 27 (25.5) |
| <i>In general, how many trials of antiepileptic drugs (AEDs) should be tried before considering the diagnosis of drug resistant epilepsy?</i> | |
| Failure of seizure control after 1 AED | 1 (0.9) |
| Failure of seizure control after 2 AEDs (correct answer) | 70 (66) |
| Failure of seizure control after ≥3 AEDs | 34 (32.1) |
| Failure of all approved AEDs | 1 (0.9) |
| I don't know | 0 (0) |
| <i>In general, how many AEDs would you try (assuming an adequate trial at an adequate dose) before referring a patient who remains drug resistant for consideration of epilepsy surgery?</i> | |
| Failure of seizure control after 1 AED | 1 (0.9) |
| Failure of seizure control after 2 AEDs (correct answer) | 53 (50) |
| Failure of seizure control after ≥3 AEDs | 49 (46) |
| Failure of all approved AEDs | 2 (1.9) |
| I don't know | 1 (0.9) |
| <i>For how long does a patient have to be drug resistant before you consider referral for epilepsy surgery evaluation?</i> | |
| As early as possible (correct answer) | 50 (47.2) |
| 3 months - 1 year | 42 (39.6) |
| 2 - 5 years | 10 (9.4) |
| > 5 years | 0 (0) |
| No one should be referred for epilepsy surgery | 0 (0) |
| I don't know | 4 (3.8) |
| <i>Patients with focal epilepsy and a normal MRI may benefit from epilepsy surgery.</i> | |
| Yes (correct answer) | 89 (84) |
| No or I don't know | 17 (16) |
| <i>Patients with generalized (non-focal) epilepsies are not candidates for epilepsy surgery.</i> | |
| Yes or I don't know | 40 (37.7) |
| No (correct answer) | 66 (62.3) |
| <i>Patients with a developmental delay are not candidates for epilepsy surgery.</i> | |
| Yes or I don't know | 15 (14.2) |
| No (correct answer) | 91 (85.8) |
| <i>Patients with psychiatric comorbidities are not candidates for epilepsy surgery.</i> | |
| Yes or I don't know | 24 (22.6) |
| No (correct answer) | 82 (77.4) |
| <i>Patients with epileptic encephalopathies are not candidates for epilepsy surgery.</i> | |
| Yes or I don't know | 34 (32.1) |
| No (correct answer) | 72 (67.9) |
| <i>Which one of the following is the most common neurological complication of epilepsy surgery?</i> | |
| Aphasia | 10 (9.4) |
| Visual field loss (correct answer) | 27 (25.5) |
| Paralysis | 11 (10.4) |
| Memory loss | 46 (43.4) |
| I don't know | 12 (11.3) |

Table 3 - Responses to the attitude section.

| Statements | n (%) |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|
| <i>How often do you discuss surgical options — when indicated — with your epilepsy patients? (positive statement)</i> | |
| 0 = Never | 3 (2.8) |
| 1 = Rarely | 14 (13.2) |
| 2 = Sometimes | 38 (35.8) |
| 3 = Very often | 28 (26.4) |
| 4 = Always | 23 (21.7) |
| <i>Epilepsy surgery is a dangerous procedure (negative statement)</i> | |
| 0 = Strongly agree | 1 (0.9) |
| 1 = Somewhat agree | 18 (17) |
| 2 = Neither agree nor disagree | 24 (22.6) |
| 3 = Somewhat disagree | 34 (32.1) |
| 4 = Strongly disagree | 29 (27.4) |
| <i>Epilepsy surgery should be viewed as a last resort for patients with epilepsy. (negative statement)</i> | |
| 0 = Strongly agree | 11 (10.4) |
| 1 = Somewhat agree | 25 (23.6) |
| 2 = Neither agree nor disagree | 12 (11.3) |
| 3 = Somewhat disagree | 33 (31.1) |
| 4 = Strongly disagree | 25 (23.6) |
| <i>If I had epilepsy that was amenable to surgical therapy, I would agree to the option of epilepsy surgery. (positive statement)</i> | |
| 4 = Strongly agree | 66 (62.3) |
| 3 = Somewhat agree | 31 (29.2) |
| 2 = Neither agree nor disagree | 7 (6.6) |
| 1 = Somewhat disagree | 1 (0.9) |
| 0 = Strongly disagree | 1 (0.9) |
| <i>If one of my relatives had epilepsy that was amenable to surgical therapy, I would encourage him/her to undergo epilepsy surgery. (positive statement)</i> | |
| 4 = Strongly agree | 68 (64.4) |
| 3 = Somewhat agree | 30 (28.3) |
| 2 = Neither agree nor disagree | 5 (4.7) |
| 1 = Somewhat disagree | 0 (0) |
| 0 = Strongly disagree | 3 (2.8) |
| <i>Epilepsy surgery is an underutilized treatment method of epilepsy. (positive statement)</i> | |
| 4 = Strongly agree | 66 (62.3) |
| 3 = Somewhat agree | 23 (21.7) |
| 2 = Neither agree nor disagree | 9 (8.5) |
| 1 = Somewhat disagree | 7 (6.6) |
| 0 = Strongly disagree | 1 (0.9) |
| <i>Specialized epilepsy centers should be available in all tertiary hospitals. (positive statement)</i> | |
| 4 = Strongly agree | 76 (71.7) |
| 3 = Somewhat agree | 19 (17.9) |
| 2 = Neither agree nor disagree | 5 (4.7) |
| 1 = Somewhat disagree | 4 (3.8) |
| 0 = Strongly disagree | 2 (1.9) |
| <i>I think epilepsy surgery is a cost-effective treatment option. (positive statement)</i> | |
| 4 = Strongly agree | 70 (66) |
| 3 = Somewhat agree | 21 (19.8) |
| 2 = Neither agree nor disagree | 8 (7.5) |
| 1 = Somewhat disagree | 5 (4.7) |
| 0 = Strongly disagree | 2 (1.9) |

scale with options of strongly agree, somewhat agree, neither agree nor disagree, somewhat disagree, and strongly disagree, with the exception of one question,

which assessed the frequency of discussing epilepsy surgery with patients. The responses were scored from 0 to 4 based on whether the attitude statement was positive or negative. Individuals who scored 25 or more




Figure 1 - The correlation between knowledge and attitude scores.

were considered to have a positive attitude. The cutoff points for sufficient or insufficient knowledge and a positive or negative attitude were calculated based on the median score in each section. The questionnaire was subsequently sent to 3 consultants and academic staff in neurology for content validation. A pilot study of 10 neurologists was conducted to assess the clarity of the questions.

Participants. We calculated a required sample size of 96 ($Z\alpha$: 1.96 for 95% confidence level; proportion: 0.5; d : 0.1). The eligibility criteria required individuals to be certified neurologists working in Saudi Arabia. We excluded neurologist who do not manage patients with epilepsy. The questionnaire link was electronically sent to 200 neurologists chosen by convenient sampling. Data were collected in January 2017. The participants provided informed consent by clicking on “agree to participate” icon in the first page of the electronic survey.

Data analysis. Data were analyzed using the Statistical Package for the Social Sciences (IBM, Armonk, NY, USA), version 23. Descriptive data are reported as mean values, standard deviations, and percentages. Categorical variables were compared using the Chi-square test or Fisher’s exact test, as appropriate. Pearson correlation was used to assess the relationships between different variables. Only differences at $p < 0.05$ were considered statistically significant.

Results. Study subjects. One hundred and six adult and pediatric neurologists met our inclusion criteria, Table 1 presents their demographics. Our sample consisted of different subspecialists, which primarily included adult general neurology (22.6%), pediatric

neurology (32.1%), epilepsy (16.0%), stroke (12.3%), neuromuscular (6.6%), neuroimmunology (3.8%), clinical neurophysiology (2.8%), movement disorders (1.9%), neurocritical care (0.9%) and neuro-cognitive disorders (0.9%). Seventy-two percent of the participants were practicing in tertiary healthcare centers. Eighty percent of the participants had at least one epilepsy center in their city, and 78% indicated that they had access to adequate expertise and resources to enable the appropriate selection of epilepsy surgical candidates. More than half of the sample (57%) saw at least 20 patients with epilepsy each month. Approximately 45% of the participants had referred less than 3 patients to the Epilepsy Monitoring Unit (EMU) for pre-surgical workup in the previous year, 33% had referred 3-10 patients, and only 22% had referred more than 10 patients. With respect to epilepsy surgery referral, 68% of the neurologists had referred less than 3 patients, 20% had referred between 3 and 10 patients and only 12% had referred more than 10 patients in the previous year. Forty percent of the participants indicated that it requires less than 3 months for an epilepsy specialist to see their patients, 38% indicated that it requires 3–6 months, and 22% indicated that it requires more than 6 months. Approximately 27% of the neurologists indicated that it requires less than 3 months for their patients to be evaluated for epilepsy surgery (i.e., EMU, neuropsychology) in the center nearest to them, 35% indicated that it requires 3–6 months, and 38% indicated that it requires more than 6 months for this process.

Knowledge. The majority of the participants (73%) stated that they are familiar with the overall content of the American Academy of Neurology clinical practice

guidelines on temporal lobe and localized neocortical resections for epilepsy, whereas 67% believed that they have sufficient knowledge regarding epilepsy surgery. The results show that only 57.5% of the neurologists had a sufficient level of knowledge regarding epilepsy surgery, with a mean score of 6.76 out of 11 and a standard deviation of 2.29. The participants' answers to the knowledge section are shown in Table 2. There was significant variability in the level of knowledge among the neurologists who obtained their last certificate of neurology from different areas ($p=0.001$): 71% of the neurologists who obtained it from North America had a sufficient level of knowledge, compared with 52% of the neurologists who obtained it from Saudi Arabia and only 10% of the neurologists who obtained it from other areas.

Neurologists with sufficient knowledge were likely to be University staff ($p=0.026$), have access to adequate expertise and resources ($p=0.043$), and refer more patients to EMU for pre-surgical work-up ($p=0.019$). A similar non-significant correlation existed with referral for epilepsy surgery. The neurologists with a high level of knowledge discussed epilepsy surgery more often with their patients ($p=0.005$). The neurologists with an insufficient level of knowledge tended to believe that surgery should be viewed as a last resort for patients with epilepsy ($p<0.001$). The neurologists with a sufficient level of knowledge toward epilepsy surgery agreed that it is an underutilized and cost-effective treatment, $p=0.01$ and $p=0.041$, respectively.

Attitude. The vast majority of the respondents discuss epilepsy surgery with their patients (97.2%); however, only 48% discuss it frequently. When asked if surgery should be viewed as a last resort for patients with epilepsy, 34% of our sample agreed (Table 3). Ninety-one percent of the neurologists indicated that they would agree to have epilepsy surgery if it was indicated for them. A majority of the neurologists (85.8%) considered epilepsy surgery a cost-effective treatment method, and 90% indicated that it should be available in all tertiary hospitals. Overall, more than half of the neurologists (52.8%) had a positive attitude toward epilepsy surgery. Neurologists who trained abroad were more likely to have a positive attitude ($p=0.02$). As shown in Figure 1, there was a significantly positive correlation ($r=0.437$, $p<0.001$) between the scores of knowledge and attitude.

Reliability of the scale. The internal consistency reliability of the overall scale was relatively acceptable (Cronbach $\alpha = 0.67$). Both the knowledge and attitude sections were consistent, with a Cronbach α of 0.65 for knowledge and 0.69 for attitude.

Discussion. This study demonstrates that approximately half of neurologists in Saudi Arabia have adequate knowledge and positive attitudes toward epilepsy surgery. Several previous studies have evaluated neurologists' knowledge of and attitudes toward epilepsy surgery. These studies have reported variable findings, many of which concur with our results.¹⁶⁻¹⁹ The response rate was 53%, which is higher than the average response rate for online surveys.²⁰ With the high response rate from both adult and pediatric neurologists and the use of a reliable instrument, we believe that this study provides a reasonable perspective on the knowledge and attitudes toward epilepsy surgery in Saudi Arabia.

Although surgery is an effective evidence-based method to treat epilepsy, a substantial proportion (42.5%) of neurologists did not have a sufficient level of knowledge regarding it. This finding may be attributed to the low number of epilepsy surgery centers in Saudi Arabia. It may also be a result of the fast development of sub-specialized neurology clinics. One study showed that 65.4% of patients relied on their neurologist or neurosurgeon as the main source of information regarding epilepsy surgery.¹³ The presence of a sufficient level of knowledge varied based on different variables. More participants who obtained their most recent certificate in neurology from North America had sufficient knowledge, this is likely due to the exposure to epilepsy surgery early in their careers as it is more developed in these countries. Individuals who work at university and other tertiary hospitals were more likely to have a sufficient level of knowledge regarding epilepsy surgery. This may be attributed to the fact that most of these hospitals are tertiary centers and provide scholarship programs that enable their neurologists to complete neurology residency training or fellowship abroad. Our study showed that individuals with adequate access to expertise and resources to enable the appropriate selection of epilepsy surgical candidates had more sufficient knowledge, which was expected because it makes them more engaged in and informed about the advances and achievements in this field. Neurologists with sufficient knowledge tended to discuss epilepsy surgery with their patients more frequently. We believe that their adequate knowledge regarding the substantial benefits of epilepsy surgery made them more comfortable proposing the surgical option to their patients. It is important to note that individuals with sufficient knowledge regarding surgery are more likely to develop positive attitudes, likely due to better understanding of indications, risks and benefits of the procedure. This relationship between knowledge and attitudes may also be because individuals with sufficient

knowledge may refer more patients for evaluation for epilepsy surgery which results in positive feedback to the referring physicians who then can appreciate the benefits of surgery on their operated patients.

Neurologist in the Kingdom of Saudi Arabia listed the health system organization, patient fears, information and motivation for patients, availability of education, and hospital organization and administration as the most important obstacles related to epilepsy surgery. Only one-third of the participants considered funding an issue. Interestingly, epilepsy surgery was not discussed frequently by 52% of the participants and not at all by 2.8%. In a previous study conducted by Hakimi et al¹⁹ the latter rate was higher at 11%. With regards to cost-effectiveness, 85.8% of our participants considered epilepsy surgery a cost-effective treatment, which is consistent with the results of a study done among Swedish neurologists (92%).¹⁸ In the present study, 34% of neurologists considered epilepsy surgery as the last resort, which may be attributed to the misunderstanding of DRE, as only 66% defined it as the failure of seizure control after 2 appropriately chosen AEDs. Among the limitations to consider is the accuracy of survey type assessments, although the content validity was assessed further validation testing of the questionnaire is required to determine its measurement properties. Participants who answered "I don't know" were considered to lack knowledge. This might not represent their real world practice, as these individuals might actually seek out knowledge when handling certain cases to determine proper disposition of patients.

In conclusion, neurologists in Saudi Arabia appear to have positive attitudes toward epilepsy surgery despite a moderate level of knowledge as measured in this study. Different variables, including the place of the last neurology certificate, type of practicing hospital, and access to expertise and resources, affected their knowledge. Adequate knowledge was positively correlated with attitude.

Acknowledgements. *We thank the neurologists who participated in this study. We acknowledge the efforts of the Neuroscience forum and Dr. Mohammed Jan in facilitating the data collection process. We also extend our gratitude to Dr. Abdullah AlRuwaite for his help in searching the literature. Finally, we would like to thank Nature Research Editing Service for English language editing.*

References

- Rajeh S, Awada A, Bademosi O, Ogunniyi A. The prevalence of epilepsy and other seizure disorders in an Arab population: a community-based study. *Seizure* 2001; 10: 410-414.
- French JA. Refractory epilepsy: clinical overview. *Epilepsia* 2007; 48: 3-7.
- Kwan P, Arzimanoglou A, Berg AT, Brodie MJ, Allen Hauser W, Mathern G, et al. Definition of drug resistant epilepsy: consensus proposal by the ad hoc Task Force of the ILAE Commission on Therapeutic Strategies. *Epilepsia* 2010; 51: 1069-1077.
- Harden CL, Maroof DA, Nikolov B, Fowler K, Sperling M, Liporace J, et al. The effect of seizure severity on quality of life in epilepsy. *Epilepsy Behav* 2007; 11: 208-211.
- Carran MA, Kohler CG, O'Connor MJ, Cloud B, Sperling MR. Marital status after epilepsy surgery. *Epilepsia* 1999; 40: 1755-1760.
- Kwan P, Schachter SC, Brodie MJ. Drug-resistant epilepsy. *N Engl J Med* 2011; 365: 919-926.
- Begley CE, Annegers JF, Lairson DR, Reynolds TF, Hauser WA. Cost of epilepsy in the United States: a model based on incidence and prognosis. *Epilepsia* 1994; 35: 1230-1243.
- Jobst BC, Cascino GD. Resective epilepsy surgery for drug-resistant focal epilepsy: a review. *JAMA* 2015; 313: 285-293.
- Engel JJ, Wiebe S, French J, Sperling M, Williamson P, Spencer D, et al. Practice parameter: temporal lobe and localized neocortical resections for epilepsy: report of the Quality Standards Subcommittee of the American Academy of Neurology, in association with the American Epilepsy Society and the American Association of Neuro. *Neurology* 2003; 60: 538-547.
- Tellez-Zenteno JF, Dhar R, Hernandez-Ronquillo L, Wiebe S. Long-term outcomes in epilepsy surgery: antiepileptic drugs, mortality, cognitive and psychosocial aspects. *Brain* 2007; 130: 334-345.
- Engel JJ. Why is there still doubt to cut it out? *Epilepsy Curr* 2013; 13: 198-204.
- Barth JH, Misra S, Aakre KM, Langlois MR, Watine J, Twomey PJ, et al. Why are clinical practice guidelines not followed? *Clin Chem Lab Med* 2016; 54: 1133-1139.
- Hrazdil C, Roberts JI, Wiebe S, Sauro K, Vautour M, Hanson A, et al. Patient perceptions and barriers to epilepsy surgery: evaluation in a large health region. *Epilepsy Behav* 2013; 28: 52-65.
- Alsemari A, Al-Otaibi F, Baz S, Althubaiti I, Aldhalaan H, MacDonald D, et al. Epilepsy Surgery Series: A Study of 502 Consecutive Patients from a Developing Country. *Epilepsy Res Treat* 2014; 2014: 286801.
- Alaqeel A, Alebdi F, Sabbagh AJ. Epilepsy: What do health-care professionals in Riyadh know? *Epilepsy Behav* 2013; 29: 234-237.
- Erba G, Moja L, Beghi E, Messina P, Pupillo E. Barriers toward epilepsy surgery. A survey among practicing neurologists. *Epilepsia* 2012; 53: 35-43.
- Roberts JI, Hrazdil C, Wiebe S, Sauro K, Vautour M, Wiebe N, et al. Neurologists' knowledge of and attitudes toward epilepsy surgery: a national survey. *Neurology* 2015; 84: 159-166.
- Kumlien E, Mattsson P. Attitudes towards epilepsy surgery: A nationwide survey among Swedish neurologists. *Seizure* 2010; 19: 253-255.
- Hakimi AS, Spanaki MV, Schuh LA, Smith BJ, Schultz L. A survey of neurologists' views on epilepsy surgery and medically refractory epilepsy. *Epilepsy Behav* 2008; 13: 96-101.
- Cunningham CT, Quan H, Hemmelgarn B, Noseworthy T, Beck CA, Dixon E, et al. Exploring physician specialist response rates to web-based surveys. *BMC Med Res Methodol* 2015; 15: 32.