



The Cochrane Library. The single most reliable source of evidence in healthcare
The Cochrane Reviews highlighted below are available from the
Cochrane Database of Systematic Reviews (www.thecochranelibrary.com)
Cochrane Reviews are regularly updated as new evidence emerges and in response to feedback,
and the Cochrane Database of Systematic Reviews should always be consulted for the most recent version of the review.

INSUFFICIENT EVIDENCE FOR THE USE OF OMEGA 3 SUPPLEMENTS IN TREATING DEPRESSION

**New research out today concludes that there is insufficient evidence for the use of taking
an Omega 3 fatty acid supplement in treating major depressive disorder.**

Omega 3 fatty acids are widely thought to be essential for good health and are naturally found in fatty fish, such as tuna, seafood and some nuts and seeds.

Omega 3 fatty acids have been widely promoted globally and are readily available, over-the-counter supplement. These supplements have hugely increased in popularity over the last decade together with a range of other supplements including ginseng, garlic, green tea, as well as vitamins, minerals and herbal products.

More recently there have been various studies that have suggested a role for Omega 3 fatty acid supplementation in treating major depressive disorder. Adults with major depressive disorders are characterized by depressed mood or a lack of pleasure in previously enjoyed activities for at least two weeks, in the absence of any physical cause, that impact on everyday life.

Figures published by the World Health Organization in 2011 estimated major depressive disorders to account for 3% of global ill health and projections for 2030 suggest an increase to 6% or 7%.

A new Cochrane review, published today in the Cochrane Library, gathered together data from 26 randomized trials involving a total of 1,458 participants. The trials investigated the impact of giving an Omega 3 fatty acid supplement in a capsule form and compared it to a dummy pill. In one study, involving 40 participants, they also investigated the impact of the same supplementation compared to an anti-depressant treatment.

The Cochrane authors found that whilst people who were given Omega 3 fatty acids reported lower symptom scores than people with the dummy pill, the effect was small and there were important limitations that undermined their confidence in the results. Their analyses showed that although similar numbers of people experienced side effects, more data would be required to understand the risks of taking Omega 3 fatty acids.

Lead author, Katherine Appleton from Bournemouth University said, “We found a small-to-modest positive effect of Omega 3 fatty acids compared to placebo, but the size of this effect is unlikely to be meaningful to people with depression, and we considered the evidence to be of low or very low quality. All studies contributing to our analyses were of direct relevance to our research question, but most of these studies are small and of low quality.”

She added, “At present, we just don’t have enough high quality evidence to determine the effects of Omega 3 fatty acids as a treatment for major depressive disorder. It’s important that people who suffer from depression are aware of this, so that they can make more informed choices about treatment.”

Full citation: Appleton KM, Sallis HM, Perry R, Ness AR, Churchill R. Omega-3 fatty acids for depression in adults. Cochrane Database of Systematic Reviews 2014, Issue 5. Art. No.: CD004692. DOI: 10.1002/14651858.CD004692.pub3.

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd., reproduced with permission.

Authorship entitlement

Excerpts from the Uniform Requirements for Manuscripts Submitted to Biomedical Journals updated November 2003.

Available from www.icmje.org

The international Committee of Medical Journal Editors has recommended the following criteria for authorship; these criteria are still appropriate for those journals that distinguish authors from other contributors.

Authorship credit should be based on 1) substantial contributions to conception and design, or acquisition of data, or analysis and interpretation of data; 2) intellectual content; and 3) final approval of the version to be published. Authors should meet conditions 1, 2, and 3.

Acquisition of funding, collection of data, or general supervision of the research group, alone, does not justify authorship.

An author should be prepared to explain the order in which authors are listed.