

What was studied in the review?

The team searched for studies that looked at giving chloroquine and hydroxychloroquine to people with COVID-19; people at risk of being exposed to the virus; and people who have been exposed to the virus.

They found 14 relevant studies: 12 studies of chloroquine or hydroxychloroquine used to treat COVID-19 in 8569 adults; two studies of hydroxychloroquine to stop COVID-19 in 3346 adults who had been exposed to the virus but had no symptoms of infection. The authors did not find any completed studies of these medicines to stop COVID-19 in people who were at risk of exposure to the virus; studies are still under way.

The studies took place in China, Brazil, Egypt, Iran, Taiwan, North America, and Europe; one study was worldwide. Some studies were partly funded by pharmaceutical companies that manufacture hydroxychloroquine.

Author Contact: Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool, UK bsingh@liverpool.ac.uk

Copyright © 2021 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd., reproduced with permission.

Physical conditions linked to psychological distress in patients with cancer

February 18, 2021 - Among patients with cancer, having additional physical comorbidities was linked with a higher risk of experiencing psychological distress. The finding comes from a Psycho-Oncology analysis of 2017 data from the National Health Survey of Spain.

The analysis included 484 patients who reported a cancer diagnosis and 484 matched controls without a history of cancer. Compared with controls, patients with cancer reported more physical comorbidities, including chronic back pain, asthma, chronic bronchitis, urinary incontinence, prostate problems, and kidney problems. They also reported higher psychological distress and were more likely to have consulted a mental healthcare professional in the past year.

Thirty percent of patients with cancer reported significant psychological distress but only 10% had consulted a professional. Each additional physical comorbidity was associated with a 9% higher odds that patients with cancer would report having high psychological distress and a 21% higher odds that they would have consulted a mental healthcare professional.

“Comorbidities often influence the choice and management of cancer treatment. These results suggest that they could also be important for patients’ mental health in the months following diagnosis,” said the first author Dafina Petrova, PhD, of the Andalusian School of Public Health, in Spain.

URL Upon Publication: <https://onlinelibrary.wiley.com/doi/10.1002/pon.5632>

Copyright © 2021 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd., reproduced with permission.