

# *Hemophilus influenzae* serotype-*b* meningitis in an adult immunocompetent patient

Hanan A. Babay, MD, KSUFelPath, Awadh R. Al-Anazy, MD, KSUFelMed, Abubakar B. Saidu, MD, Abdelmageed M. Kambal, MD, FRCPath.

---

## ABSTRACT

We present a case of meningitis caused by *Hemophilus influenzae type-b* in an immunocompetent 41-year-old Saudi lady. The patient was successfully treated with Ceftriaxone for 10 days. A review of *Hemophilus influenzae* meningitis in adults and the impact of conjugated vaccine on the epidemiology of the disease are given.

Neurosciences 2003; Vol. 8 (3): 188-190

---

*Hemophilus influenzae type-b (Hib)* is the leading cause of invasive infections such as meningitis and septicemia in children younger than 5 years.<sup>1</sup> Conjugative vaccines have had a dramatic effect on the epidemiology of the disease in this age group.<sup>2-4</sup> *Hemophilus influenzae type-b* has been recognized as a pathogen in adults particularly those with underlying illnesses.<sup>5-7</sup> Other reports suggest that the increasing incidence in such adults may be due to failure of natural acquisition of immunity to *Hib* or waning immunity with aging.<sup>8</sup> In addition, non-typable strains of *Hemophilus influenzae (Hi)* have also been isolated from adults with underlying illnesses with infections such as meningitis, pneumonia and septicemia.<sup>6,7,9</sup> On the contrary, there is a report from Alaskan natives indicating a decrease of *Hib* infections among adults after introducing the *Hib* conjugate vaccine program.<sup>10</sup> We report a case of meningitis caused by *Hib* in an apparently immunocompetent adult.

**Case Report.** A 41-year-old Saudi female patient was admitted to a medical ward with a 3 day history of

fever, chills and rigors associated with headache and vomiting. A day after onset of fever, she complained of right flank pain. She denied any further symptoms suggestive of genitourinary disorder. She had no past history of note, except for nasal polypectomy 10 months earlier, which was followed by episodes of cerebrospinal fluid (CSF) leak. Physical examination revealed an ill-looking patient, pyrexial, (38.3°C), conscious and alert with no signs of meningeal irritation. Abdomen was tender at both flanks and right hypochondrium with no palpable masses or organomegaly. Initial investigations: white blood cell count of  $21.5 \times 10^9/L$  with 91% neutrophils, erythrocyte sedimentation rate 122 mm/hour; other parameters were within normal. A midstream urine microscopy revealed 25 white blood cells/mm<sup>3</sup> and 190 red blood cells/mm<sup>3</sup>. A working diagnosis of acute pyelonephritis was made. Other investigations to rule out perinephric or hepatic abscess were made. She was put on ampicillin and gentamicin. However the patient did not show clinical response and remained febrile 24 hours after starting antibiotics. Abdomen ultrasound was unremarkable. Gram stain of

---

From the Department of Pathology/Microbiology (Babay, Kambal), Department of Medicine/Infectious Diseases (Al-Anazy, Saidu), King Khalid University Hospital, Riyadh, Kingdom of Saudi Arabia.

Received 20th November 2002. Accepted for publication in final form 15th March 2003.

Address correspondence and reprint request to: Dr. Hanan A. Babay, Department of Pathology/Microbiology (32) King Khalid University Hospital, PO Box 2925, Riyadh 11461, Kingdom of Saudi Arabia. Tel. +966 (1) 4672457. Fax. +966 (1) 4672462. E-mail: hahabib@ksu.edu.sa

positive blood culture [BacT-Alert 3D, Organon Teknika, United States of America, (USA)] showed Gram-negative coccobacilli. At this stage of her condition, the infectious diseases team was consulted. The patient was seriously ill with high fever (39°C), stiff neck, positive Kernig's sign, mentally alert and had normal fundoscopy. Clinical diagnosis of acute pyogenic meningitis was made and antibiotics changed to intra-venous ceftriaxone 4g/day. An urgent brain computerized tomography scan was carried out which did not contraindicate lumbar puncture. It showed no evidence of CSF leak or other abnormalities. Subsequent CSF examination revealed a turbid CSF with white blood cells count of 1200/mm<sup>3</sup> with 75% polymorphs and 25% lymphocytes, red blood cells were 100/mm<sup>3</sup>, glucose=0.4 mmol/L (2.5-4 mmol/L) and protein=4.4 g/L (0.15-0.4 g/L). Gram stain showed gram negative coccobacilli. Latex agglutination of CSF was positive for *Hib* (Wellcogen Bacterial antigen Kit, Murex biotech Limited, United Kingdom). The isolate from blood culture was identified as -lactamase producing *Hib* susceptible to cefuroxime, ceftriaxone, cefotaxime, gentamicin, chloramphenicol, ciprofloxacin and imipenem. The CSF culture was however negative. Immunoglobulin study was normal. Other investigations were all normal. The patient improved within 48 hours of starting ceftriaxone. A repeat lumbar puncture before discharge revealed Xanthochromic but clear CSF. She was discharged after 10 days of antibiotic course. The patient was seen 6 weeks after discharge where she was normal with no sequel.

**Discussion.** Adult cases of *Hib* meningitis are infrequent and often follow remote head trauma; prior neurosurgery or upper respiratory tract infections.<sup>5,7,11</sup> In our patient the meningitis may be associated with the previous nasal polypectomy and CSF leak. This association has also previously been reported.<sup>5,7,12</sup> Most infectious agents were suspected to reach the central nervous system by hematogenous spread, or by retrograde propagation of infected thrombi, or occasionally through bone but not through olfactory nerve.<sup>13</sup> Clinical features of meningitis in adults are similar to those in children.<sup>5,7,11</sup> However in our case only fever and headache were initially present. The neck stiffness followed later. Some studies report neck stiffness in most cases while other studies do not.<sup>11,14</sup> Our patient was conscious and had no seizures. This in contrast to the findings of Sigurdardottir et al<sup>7</sup> in which 60% of their patients had alteration in consciousness and 11% had alteration in seizures.

In KSA, the incidence of *Hib* disease dramatically decreased after the implementation of routine infant vaccination with conjugated *Hib* vaccine.<sup>2,15</sup> This was documented in one study by increased serum antibody level in response to the vaccine.<sup>15</sup> In the USA, the impact of large scale vaccination of children made *Hib* meningitis a disease of adults rather than infants and

young children.<sup>16</sup> Non-type *b-Hi* pathogen may have increased due to less competition from *Hib* in colonizing the nasopharynx.<sup>3,8,10,17,18</sup> It is too early to evaluate the effect of *Hib* vaccine in the epidemiology of invasive diseases in the KSA. The vaccine use became mandatory only recently, although since its approval by the food and drug administration, it has been available in certain private children clinics throughout the KSA since 1990.

In conclusion, meningitis and other invasive diseases due to *Hi* can occur in adult patients particularly those with underlying diseases as well as those with competent immune system. A population based surveillance program is recommended to study diseases due to *Hib* (and other serotypes) among adults in the KSA.

## References

1. Dajani AS, Asmar BI, Thirumoorthi MC. Systemic Haemophilus influenzae disease: an overview. *J Pediatr* 1979; 94: 355-364.
2. Al-Muneef M, Al-Shaalan M, Memish M, Alalola S. Bacterial meningitis in Saudi Arabia: the impact of Haemophilus influenzae type b vaccination. *J Chemother* 2001; 13 Suppl 1: 34-39.
3. Muhlemann K, Alexander ER, Pepe M, Weiss NS, Schopfer K and the Swiss Haemophilus influenzae study group. Invasive Haemophilus influenzae disease and epiglottitis among Swiss children from 1980 to 1993: Evidence of herd immunity among older age groups. *Scand J Infect Dis* 1996; 28: 265-268.
4. Centers for Disease Control: Progress towards elimination of Haemophilus influenzae type b disease among infants and children - United States 1987-1993. *MMWR Morb Mortal Wkly Rep* 1992; 43: 144-148.
5. McGee ZA, Baringer JR. Acute meningitis. In: Mandell GL, Douglas RG, Benett JE, editors. Principles and Practice of infectious diseases. 3rd ed. New York (NY): Churchill Livingstone; 1990. p. 741-755.
6. Wallace Jr RJ, Musher DM, Septimus EJ, McGowen Jr JE, Quinones FJ, Wiss K et al. Haemophilus influenzae infections in adults: characterization of strains by serotypes, biotypes and β-lactamase production. *J Infect Dis* 1981; 144: 101-106.
7. Sigurdardottir B, Bjornsson OM, Jonsdottir KE, Erlendsdottir H, Gudmundsson S. Acute bacterial meningitis in adults. A 20 year overview. *Arch Intern Med* 1997; 157: 425-430.
8. Holmes SJ, Lucas AH, Osterholm MT, Froeschle JE, Granoff DM. Immunoglobulin deficiency and idiotype expression in children developing Haemophilus influenzae type b disease after vaccination with conjugate vaccine. The Collaborative Study Group. *JAMA* 1991; 266: 1960-1965.
9. Hugosson S, Silfverdal SA, Garpenholt O, Esbjurner E, Lindquist BO, Vikerfor T et al. Invasive Haemophilus influenzae diseases: epidemiology and clinical spectrum before large-scale H. influenzae type b vaccination. *Scand J Infect Dis* 1995; 27: 63-67.
10. Perdue DG, Bulkow L, Gellin BG, Davidson M, Petersen KM, Singleton RJ et al. Invasive Haemophilus influenzae disease in Alaskan residents aged 10 years and older before and after infant vaccination programs. *JAMA* 2000; 283: 3089-3094.
11. Durand ML, Calderwood SB, Weber DJ, Miller SI, Southwick FS, Caviness VS et al. Acute bacterial meningitis in adults. A review of 493 episodes. *N Engl J Med* 1993; 328: 21-28.
12. Crawford C, Kennedy N, Weir WR. Cerebrospinal fluid rhinorrhoea and Haemophilus influenzae meningitis 37 years after head injury. *J Infect* 1994; 28: 93-97.
13. Greenlee JE. Anatomic consideration in central nervous system infections. In: Mandell GL, Douglas RG, Benett JE editors. Principles and practice of infectious diseases. 3rd ed. New York (NY): Churchill Livingstone; 1990. p. 732-741.

14. Carpenter RR, Petersdorf RG. The clinical spectrum of bacterial meningitis. *Am J Med* 1962; 33: 262-275.
15. Frayha HH, Shaheen H, Ganelin RS, Kattan HA, Tufenkeji HT, Krotzer S et al. Clinical and serologic responses of Saudi children to Haemophilus influenzae type b capsular polysaccharide diphtheria toxoid conjugate vaccine. *Annals of Saudi Medicine* 1993; 13: 215-221.
16. Schuchat A, Robinson K, Wenger JD, Harrison LH, Farley M, Reingold, et al. Bacterial meningitis in the United States in 1995. *N Engl J Med* 1997; 337: 970-976.
17. Urwin G, Krohn JA, Robinson KD, Wenger JD, Farley MM, and the Haemophilus influenzae study group. Invasive disease due to Haemophilus influenzae serotype f. Clinical and epidemiologic characteristics in the H. influenzae serotype b vaccine era. *Clin Infect Dis* 1996; 22: 1069-1076.
18. Mohle-Boetani J, Shuuchat A, Plikaytis BD, Smith DJ, Broome CV. Carriage of Haemophilus influenzae type b in children after widespread vaccination with conjugate Haemophilus influenzae type b vaccines. *Pediatr Infect Dis J* 1993; 12: 589-593.